

The page features a decorative design with three blue circles of varying sizes, each composed of concentric rings in different shades of blue. These circles are positioned in the upper right and lower right areas. Two thin blue lines intersect to form a large 'V' shape that frames the text on the left side of the page.

GRASSROOTS ACADEMY GATE-BIOTECH 2001

Correspondence Available

Grassroots is a premier GATE-BIOTECH coaching at New Delhi, with the best results including Rank 49, 71, 313 in GATE 2010 and Rank 77 in GATE 2011.

www.grassrootsacademy.in

Ph.9891602060

GATE BIOTECHNOLOGY – 2001

1. The limiting factor in the production of large quantities of ethanol as bio-fuel is
 - (A) Lack of a balanced medium
 - (B) Ethanol to toxicity to cells
 - (C) Expensive downstream processing steps
 - (D) Only (B) and (C) of the above

2. Gel retardation assay is routinely used to monitors interaction between
 - (A) Proteins
 - (B) Drug and nucleic acid
 - (C) Nucleic acid
 - (D) Protein and nucleic acid

3. Efficient expression of a heterologous protein product is influenced by
 - (A) Transcriptional efficiency
 - (B) Copy number of the plasmid
 - (C) Codon bias
 - (D) All of the above

4. The fundamental feature of the genetic code which allows the expression of a protein in any host is its
 - (A) Triplet nature
 - (B) Universality
 - (C) Degeneracy
 - (D) Redundancy

5. When thymidine synthesis is inhibited eukaryotic cells will be arrested at
 - (A) End of S phase
 - (B) End of M phase
 - (C) G1/S Interphase
 - (D) G2/ M Interphase

6. To PCR amplify the sequence
 - (A) ATCTTCTACG
 - (B) AAGCTTGCGG
 - (C) TAGAAGATGC
 - (D) TTCGAACGCC

The required primers are

 - (A) ATCTTCTA and CGAACGCC
 - (B) ATCTTCTA and CCGCAAGC
 - (C) TAGAAGAT and CGAACGCC
 - (D) TAGAAGAT and CCGCAAGC

7. The major groove of DNA is lined by
 - (A) N3 of purine and N1 of pyrimidine
 - (B) N7 of purine and O2 of pyrimidine
 - (C) N7 of purine and C4 of pyrimidine
 - (D) None of the above

8. The heterozygosity of any locus can be ascertamed by
 - (A) RFLP analysis
 - (B) SNPs
 - (C) FISH analysis
 - (D) Either RFLP analysis of SNPs

9. BAC which can be used to clone large DNA fragments, is derived from
 - (A) colE plasmid
 - (B) F plasmid
 - (C) 2 μ
 - (D) Mu phage

10. Antibody diversity is generated by
 - (A) Protein splicing
 - (B) Somatic mutation
 - (C) Allelic exclusion
 - (D) Inter-chromosomal recombination



11. Resistance to herbicide chlorsulfuron in plants is due to a change in
(A) Glutamine synthetase (B) Threonine deaminase
(C) Acetolactate synthase (D) DNA polymerase
12. A heterologous protein for its expression in the milk of transgenic animal should be under the control of the promoter of the gene coding for
(A) β -Globin (B) β -Lactoglobulin
(C) Preproinsulin (D) *lacZ*
13. To isolate a gene coding for glucagons, the cDNA library has to be constructed using mRNA isolated from
(A) Intestine (B) Pancreas
(C) Pituitary (D) Brain
14. If the fractional recovery at each step of unit operation is 0.8, the recovery after 4 steps will be
(A) 0.24 (B) 3.23
(C) 0.41 (D) 0.82
15. The factor (s) likely to increase the rate of reaction catalyzed by a surface immobilized enzyme is /are
(A) Increased agitation of the bulk liquid containing the substrate
(B) Continued replacement of the bulk liquid containing the substrate
(C) Increased concentration of the substrate in the bulk liquid
(D) All of the above
16. Which of the following cases are likely to lead to faster rate of catalysis by an enzyme immobilized on a negatively charged support?
(A) A positively charged substrate and a negatively charged product
(B) A negatively charged substrate and a positively charged product
(C) A positively charged substrate and a positively charged product
(D) None of the above
17. The essential component of Ti plasmid required for integration into plant genome is
(A) Origin of replication (B) Tumor inducing gene
(C) Nopaline utilization gene (D) All of the above
18. Positional cloning approach exploits information
(A) On the location of the gene in the genome
(B) On the status of its expression
(C) About the position of promoter relative to MCS
(D) About the position of the restriction sites
19. Hormone pairs required for a callus to differentiate are
(A) Auxin and cytokinin (B) Auxin and gibberellin
(C) Ethylene and gibberellin (D) Cytokinin and gibberellin
20. Reverse genetics means
(A) Finding the function of a ORF
(B) Finding the gene responsible for a trait
(C) RNA dependent DNA synthesis
(D) Converting somatic cell to a germ cell
21. Which of the following statements applies to the operation of a fed-batch process?
(A) The volume of the culture increases with time
(B) it helps controlling repressive effects of the nutrient being fed
(C) It eliminates the need for oxygen supply
(D) Only (A) and (B) of the above



22. A gene can not be isolated from a human genomic DNA library by functional complementation in *E. coli* because of
- (A) None functional promoter
 - (B) The absence of splicing machinery
 - (C) Coupled transcription and translation
 - (D) Codon bias
23. In large-scale fermentation the preferred method of sterilization is
- (A) Chemicals
 - (B) Radiation
 - (C) Filtration
 - (D) Heat
24. Embryo rescue is a useful technique to
- (A) Grow / generate hybrids between different plant species
 - (B) Complete the growth of embryos susceptible to defects in seed development
 - (C) Break the dormancy of seeds
 - (D) All of the above
25. Which of the following is not true of aerobic digestion?
- (A) It generates most sludge
 - (B) It generally incurs higher running cost
 - (C) It may generate a usable fuel
 - (D) Requires a shorter residence time

26. Match the products in **Column A** with their corresponding organisms in **Column B**

Column A

- (A) *Aspergillus niger*
- (B) *Saccharomyces cerevisiae*
- (C) *Penicillium chrysogenum*
- (D) *Lactobacillus casei*
- (E) *Corynebacterium glutamicum*

Column B

- 1. Lysine
- 2. Citric acid
- 3. Acetone / butanol
- 4. Vitamin B₁₂
- 5. Erythromycin
- 6. β- Lactam
- 7. Diacetyl
- 8. Ethanol

27. Match the unit operations in **Column A** with the most **appropriate** recovery stage in **Column B**

Column A

- (A) Drying
- (B) Sedimentation
- (C) Membranes
- (D) Cell disruption
- (E) Chromatography

Column B

- 1. Pretreatment
- 2. Purification
- 3. Formulation
- 4. Solid/ liquid separations
- 5. Solid / solid separation
- 6. Concentration

28. (a) A chemostat is operating at state at a dilution rate of 0.1 hr^{-1} and a limiting nutrient concentration of $10 \mu\text{M}$. If the μ_{max} for the organism is 0.5 hr^{-1} calculate the Monod constant for the nutrient.
(b) Why asparagine is used in anticancer therapy?
29. (a) Write the reaction catalyzed by penicillin G acylase
(b) Name any two techniques by which penicillin G acylase may be immortalized
(c) Why are mammalian cells cultured in CO₂ incubators?
(d) Mention an important post translational modification absent in prokaryotes making them unsuitable hosts for expressing human genes.



30. (a) The sequences at the cloning site of three vectors are given below. The *Bam*H I (GGATCC) and *Hind* III (AAGCTT) sites are underlined. Only the sequences around the restriction site are shown. The symbol "_____" indicates rest of the sequence.

Vector 1: ____ Promoter ...ATGGGTCGCGGATCCGGCTGC...AAGCTT ____

Vector 2: ____ Promoter ...ATGGGTCGGGATCCGGCTGCT...AAGCTT ____

Vector3: ____ Promoter...ATGGGTCGATCCGGCTGCTA...AAGCTT ____

Which one of the above three vectors is appropriate to clone the following ORF

_____ ATGCCCAACACCCGATCCCG... TAAAAGCTT _____

For expression? Give the reaction in one sentence.

- (b) Draw the restriction map of the plasmid given the following data (the gel pattern shown below is not to scale). The size of each DNA fragment (in kb) is indicated next to it.

<i>eco</i> R 1	<i>Sal</i> I	<i>Hind</i> III	<i>Eco</i> R I & <i>Hind</i> III	<i>Sal</i> I & <i>Hind</i> III	<i>Eco</i> R 1 & <i>Sal</i> I
5.4__	5.4__				
					3.6__
		2.1__	2.1__		
		1.9__		1.9__	1.8__
		1.4__	1.4__	1.4__	
			1.3__	1.2__	
				0.9__	
			0.6__		

