



GRASSROOTS ACADEMY **GATE- BIOTECH-2011**

Correspondence Available

Grassroots is a premier GATE-BIOTECH Coaching at New Delhi, with the Best results including Rank 49, 71, 313 in GATE 2010 and Rank 77 in GATE 2011.

www.grassrootsacademy.in

BIOTECHNOLOGY - 2011

1 – 25 carry one mark each.

- Embryonic stem cells are derived from
(A) Fertilized embryo (B) unfertilized embryo
(C) Sperm (D) kidney
- Members of the antibody protein family that have common structural features are collectively known as
(A) haptens (B) allergens
(C) antigens (D) immunoglobulins
- Apoptosis is characterized by
(A) Necrosis (B) Programmed cell death
(C) Membrane leaky syndrome (D) Cell cycle arrest process
- Yeast artificial chromosomes (YAC's) are used for cloning
(A) Large segments of DNA (B) mRNA
(C) Bacterial DNA (D) Yeast DNA
- The product commercially produced by animal cell culture is
(A) Insulin (B) tissue plasminogen activator
(C) Interferon (D) Hepatitis B vaccine
- An alternative to glycolysis pathway is
(A) Glyoxylate pathway (B) Pentose phosphate pathway
(C) Citric acid cycle (D) Gluconeogenesis
- A cell in G_1 of interphase has 12 chromosomes. How many chromatids will be found per cell during metaphase II of meiosis?
(A) 6 (B) 12
(C) 18 (D) 24
- Diploid *Drosophila* has eight chromosomes. Which one of the following terms should NOT be used to describe *Drosophila* with sixteen numbers of chromosomes?
(A) Polyploid (B) Aneuploid
(C) Euploid (D) Tetraploid
- Hydrated synthetic seeds which are produced by ion exchange reaction involve mixing the somatic embryos in a solution of
(A) Sodium alginate and dropping it in a solution of calcium nitrate
(B) Calcium alginate and dropping it in a solution of sodium nitrate
(C) Calcium alginate and dropping it in a solution of ammonium nitrate

(C) α_1 and β_1 domains only

(D) α_1 and β_2 domains only

25. The protein in eukaryotes which is subjected to degradation undergoes

(A) Phosphorylation

(B) Carboxylation

(C) Ubiquitination

(D) Methylation

26 – 55 carry two marks each.

26. Match the viruses in **Group I** with their host cell receptors in **Group II**.

Group I

P. Hepatitis A virus

Q. Human immunodeficiency virus

R. Rabies virus

S. Herpes simplex virus type I

Group II

1. Heparan sulphate

2. Acetylcholine receptor

3. CD4 protein

4. Alpha-2 macroglobulin

(A) P – 1, Q – 3, R – 2, S – 4

(B) P – 3, Q – 4, R – 1, S – 2

(C) P – 4, Q – 3, R – 2, S – 1

(D) P – 2, Q – 3, R – 1, S – 4

27. Match the microbial growth characteristics in **Group I** with the corresponding features in **Group II**.

Group I

P. Growth associated product formation

Q. Non growth associated product formation

R. Product inhibition

S. Substrate inhibition

Group II

1. Specific growth rate decreases increasing product concentration

2. Specific product formation rate is constant

3. Specific product formation rate is proportional to specific growth rate

4. Specific growth rate decreases With increasing substrate -
- concentration

(A) P-1, Q-2, R-4, S-3

(B) P-3, Q-2, R-1, S-4

(C) P-2, Q-1, R-3, S-4

(D) P-2, Q-3, R-4, S-1

28. Match the items in **Group I** with **Group II**.

Group I

P. Circular dichroism

Q. X-ray crystallography

R. Freeze-drying

Group II

1. Concentration

2. Sedimentation coefficient

3. Secondary structure determination

S. Ultracentrifugation

4. Tertiary structure determination

(A) P-4, Q-1, R-2, S-3

(B) P-1 Q-4, R-3, S-2

(C) P-2, Q-3, R-4, S-1

(D) P-3, Q-4, R-1, S-2

29. Match the products in **Group I** with their respective organisms in **Group II**.

Group I

P. Glycerol

Q. Glutamic acid

R. Curdlan

S. Amphoterin B

Group II

1. *Corynebacterium glutamicum*

2. *Alcaligenes faecalis*

3. *Dunaliella salina*

4. *Streptomyces nodosus*

(A) P-2, Q-1, R-3, S-4

(B) P-4 Q-2, R-1, S-3

(C) P-3, Q-1, R-2, S-4

(D) P-2, Q-1, R-4, S-3

30. Determine the correctness or otherwise of the following **Assertion (a)** and the **Reason (r)**.

Assertion: I_gM is found in serum as a pentameric protein consisting of five I_gM monomers.

Reason: The pentameric form of I_gM is due to cross-linking of I_gM monomers via peptide bond.

(A) Both (a) and (r) are true and (r) is the correct reason for (a)

(B) Both (a) and (r) are true but (r) is not the correct reason for (a)

(C) (a) is true but (r) is false

(D) (a) is false but (r) is true

31. Determine the correctness or otherwise of the following **Assertion (a)** and the **Reason (r)**.

Assertion: N-methyl-N' -nitro-N-nitrosoguanidine (NTG) is an effective chemical mutagen.

Reason: Mutations induced by NTG mainly are the $GC \rightarrow AT$ transitions.

(A) Both (a) and (r) are true and (r) is the correct reason for (a)

(B) Both (a) and (r) are true but (r) is not the correct reason for (a)

(C) (a) is true but (r) false

(D) (a) is false but (r) is true

32. Determine the correctness of the following statements

I. Enhancer sequences are those DNA sequences that are involved in increasing, the rate of DNA replication.

II. Enhancer sequences work by binding with eukaryotic gene activator factors.

- (A) Only I is true
 (B) Only II is true
 (C) Both I and II are true
 (D) Both I and II are false

33. In a well aerated and agitated microbial culture, the 'supply' of oxygen is equal to 'demand' (uptake) of the growing culture. The $K_L a$ for such a system will be ($K_L a$ = volumetric mass transfer coefficient, C^* = dissolved oxygen concentration in liquid in equilibrium with gaseous oxygen, C = instantaneous value of dissolved oxygen concentration, ' r ' = specific oxygen uptake rate per unit weight of cells, X = dry weight of the cells per unit volume.

- (A) $(r X) / (C^* - C)$
 (B) $(r) / X (C^* - C)$
 (C) $(C) (C^* - C) / (r X)$
 (D) $(X) / R (C^* - C)$

34. Structured William's model

- P. can describe the changes in intracellular components of the cell during growth
 Q. can not describe the death phase of the cells
 R. can describe the variation of size of the cells in the different phases of growth
 S. can not describe the lag period of growth

Which one of the following is **CORRECT**?

- (A) P, Q and S only
 (B) P, Q and R only
 (C) Q, R and S only
 (D) P, Q and S only

35. Match items in **Group I** with **Group II**.

Group I

- P. Glycolytic pathway
 Q. Eukaryotic oxidative metabolism
 R. Glyoxylate cycle
 S. Calvin cycle

- (A) P-1, Q-2, R-3, S-4
 (C) P-4, Q-3, R-2, S-1

Group II

1. Chloroplast
 2. Glyoxysomes
 3. Mitochondria
 4. Cytosol

- (B) P-2 Q-3, R-4, S-1
 (D) P-3, Q-4, R-1, S-2

36. Match items in **Group I** with **Group II**.

Group I

- P. Alzheimer's disease
 Q. Mde cow disease
 R. Sickle cell anaemia
 S. Swine flu

Group II

1. H1 N1
 2. Hemoglobin
 3. Prions
 4. Amyloid



(A) P-4, Q-3, R-2, S-1
(C) P-2, Q-1, R-4, S-3

(B) P-3 Q-4, R-1, S-2
(D) P-1, Q-2, R-3, S-4

37. Determine the correctness or otherwise of the following **Assertion** (a) and the **Reason** (r).

Assertion: The elucidation of ribosome structure helps in the development of new generation drugs.

Reason: The high resolution of macromolecular structure has enabled in structure-based drug design.

(A) Both (a) and (r) are true and (r) is the correct reason for (a)
(B) Both (a) and (r) are true but (r) is not the correct reason for (a)
(C) (a) is true but (r) is false
(D) (a) is false but (r) is true

38. Determine the correctness or otherwise of the following **Assertion** (a) and the **Reason** (r).

Assertion: A very low amount of inhibitor can act as an activator for allosteric enzymes.

Reason: Allosteric enzymes follow Michaelis-Menten kinetics.

(A) Both (a) and (r) are true and (r) is the correct reason for (a)
(B) Both (a) and (r) are true but (r) is not the correct reason for (a)
(C) (a) is true but (r) is false
(D) (a) is false but (r) is true

39. Match the terms in **Group I** with their associated functions in **Group II**.

Group I

P. Shine-Dalgarno sequences
Q. Leucine zipper
R. Aminoacyl tRNA synthetase
S. RNA interference (RNAi)

Group II

1. Aminoacylation of tRNA
2. Gene silencing
3. Ribosome binding and facilitation of
-translation initiation
4. Transcription factors

(A) P-3, Q-4, R-1, S-2
(C) P-2, Q-3, R-1, S-4

(B) P-4 Q-3, R-2, S-1
(D) P-3, Q-2, R-4, S-1

40. Protein-protein interactions are studied by

P. DNA foot printing
Q. Yeast two hybrid system
R. Ligase chain reaction

S. Mass spectrometry

- (A) P and S only
(B) Q and S only
(C) P and R only
(D) Q and R only

41. Determine the correctness or otherwise of the following **Assertion** (a) and **Reason** (r)

Assertion: Isopropylthiogalactoside (PITG) is a gratuitous inducer of lactose operon.

Reason: Gratuitous inducers are chemical analogs which behave like natural inducer but they do not serve as substrate for the enzymes that are subsequently synthesized.

- (A) Both (a) and (r) are true and (r) is the correct reason for (a)
(B) Both (a) and (r) are true but (r) is not the correct reason for (a)
(C) (a) is true but (r) is false
(D) (a) is false but (r) is true

42. Determine the correctness or otherwise of the following **Assertion** (a) and the **Reason** (r).

Assertion: In synchronous culture, majority of the cells move to next phase of the cell cycle simultaneously.

Reason: Synchronous culture could be obtained by starving cells for essential nutrient components.

- (A) Both (a) and (r) are true and (r) is the correct reason for (a)
(B) Both (a) and (r) are true but (r) is not the correct reason for (a)
(C) (a) is true but (r) is false
(D) (a) is false but (r) is true

43. Which of the following characteristics with respect to bacterial DNA polymerase III are **TRUE**?

- P. Initiation of chain synthesis
Q. 5' - 3' polymerization
R. 5' - 5' exonuclease activity
S. 5' - 3' exonuclease activity

- (A) P and Q only
(B) Q and R only
(C) Q and S only
(D) P and S only

44. Maximum specific growth rate (μ_{max}) of a microorganism is calculated by taking the ($\ln = \log_e$, X = biomass, t = time)

- (A) slope of $\ln X$ vs t of the growth cycle
(B) slope of $\ln X$ vs t during the exponential growth phase

- (C) slope of X vs t
 (D) slope of X vs t during the exponential phase of growth

45. Identify the **CORRECT** statements

- P. 5' AND 3' ends of the transcripts can be mapped by utilizing polymerase chain reaction.
 Q. S_1 nuclease can cleave the DNA strand of a DNA-RNA hybrid
 R. T_4 polynucleotide kinase is used for labeling 3' end of DNA
 S. Baculovirus (*Autographa californica*) can be used as an insect expression vector

- (A) P and Q only
 (B) R and S only
 (C) P and S only
 (D) Q and R only

46. Value of the determinant mentioned below is

$$\begin{vmatrix} 1 & 0 & -1 & 0 \\ 4 & 7 & 0 & 2 \\ 1 & 1 & -1 & 1 \\ 2 & 0 & 2 & 1 \end{vmatrix}$$

- (A) 24
 (B) -30
 (C) -24
 (D) -10

47. HAT (hypoxanthine, aminopterin and thymidine) is used for selecting the hybridomas based on the following

- I Only hybridoma will grow since it inherited the HGPRT genes from B-cells and can synthesize DNA from hypoxanthine.
 II. Myeloma cells will not grow in cultures since *de novo* synthesis is blocked by aminopterin and due to the lack of HGPRT enzyme.

- (A) Only I is true
 (B) Only II is true
 (C) Both I and II are true
 (D) I is true and II is false

Common data questions 48 and 49

Red-green colour blindness is inherited as a recessive X-linked trait.

48. What will be the probability of having the colour-blind son to a woman with phenotypically normal parents and a colour-blind brother, and married to a normal man? (assume that she has no previous children)
 (A) 100%
 (B) 50%

- (C) 25% (D) 12.5%

49. What will be the probability of having the colour-blind daughter to a phenotypically normal woman, who already had one colour-blind son, and is married to a colour-blind man?
- (A) 75% (B) 50%
(C) 25% (D) 15%

Common data for Questions 50 and 51

A microorganism grows in a continuous 'chemostat' culture of 60 m³ working volume with sucrose as the growth limiting nutrient at dilution rate, $D = 0.55 \text{ h}^{-1}$. The steady state biomass concentration is 4.5 Kg dry biomass m⁻³ and the residual sucrose concentration is 2.0 Kg m⁻³. The sucrose concentration in the incoming feed medium is 10.0 Kg m⁻³.

50. What would be the yield $Y_{x/s}$ (Kg biomass/Kg substrate)?
- (A) 0.562 (B) 0.462
(C) 0.362 (D) 0.162
51. What would be the sucrose concentration in the input feed for the output to be 45 Kg biomass h⁻¹?
- (A) 3.225 Kg m⁻³ (B) 4.425 Kg m⁻³
(C) 5.115 Kg m⁻³ (D) 6.525 Kg m⁻³

Linked Answer Questions

Statement for Linked Answer Questions 52 and 53.

The abdomen length (in millimeters) was measured in 15 male fruit flies, and the following data were obtained: 1.9, 2.4, 2.1, 2.0, 2.2, 2.4, 1.7, 1.8, 2.0, 2.0, 2.3, 2.1, 1.6, 2.3, and 2.2.

52. Variance (V_x) for this population of fruit flies as calculated from the above data shall be
- (A) 0.85 (B) 0.25
(C) 0.061 (D) 0.08
53. The value of standard deviation (SD) will be
- (A) 0.061 (B) 0.25
(C) 0.61 (D) 0.85

Statement for Linked Answer Questions 54 and 55:



A 200 μl of polymerase chain reaction has 100 template DNA molecules and the reaction was performed for 10 cycles.

54. How many molecules of amplicons will be generated?
(A) 1.014×10^4 (B) 1.024×10^5
(C) 2.048×10^4 (D) 2.048×10^5
55. How many molecules of amplicons will be present in 0.1 μl of reaction?
(A) 102.4 (B) 1024
(C) 51.2 (D) 512

General Aptitude (GA) Questions.

Q. 56 – 60 carry one mark each

56. Which of the following options is the closest in the meaning to the word below:
Inexplicable
(A) Incomprehensible (B) Indelible
(C) Inextricable (D) Infallible
57. Choose the word from the options given below that is most nearly opposite in meaning to the given word: **Amalgamate**
(A) Merge (B) Split
(C) Collect (D) Separate
58. Choose the most appropriate word from the options given below to complete the following sentence.
If you are trying to make a strong impression on your audience, you cannot do so by being understated, tentative or _____
(A) Hyperbolic (B) Restrained
(C) Argumentative (D) Indifferent
59. Choose the most appropriate word(s) from the options given below to complete the following sentence.
I contemplated _____ Singapore for my vacation but decided against it.
(A) To visit (B) having to visit
(C) Visiting (D) for a visit
60. If $\text{Log}(P) = (1/2) \text{Log}(Q) = (1/3) \text{Log}(R)$, then which of the following options is **TRUE**?
(A) $P^2 = Q^3 R^2$ (B) $Q^2 = PR$

(C) $Q^2 = R^3 P$

(D) $R = P^2 Q^2$

Q. 61 – 65 carry two marks each

61. Few school curricula include a unit on how to deal with bereavement and grief, and yet all students at some point in their lives suffer from losses through death and parting.
Based on the above passage which topic would not be included in a unit on bereavement?
- (A) How to write a letter of condolence
(B) What emotional stages are passed through in the healing process?
(C) What the leading causes of death are
(D) How to give support to a grieving friend
62. A container originally contains 10 liters of pure spirit. From this container 1 litre of spirit is replaced with 1 litre of water. Subsequently, 1 litre of the mixture is again replaced with 1 litre of water and this process is repeated one more time. How much spirit is now left in the container?
- (A) 7.58 litres (B) 7.84 litres
(C) 7 litres (D) 7.29 litres
63. A transporter receives the same number of orders each day. Currently, he has some pending orders (backlog) to be shipped. If he uses 7 trucks, then at the end of the 4th day he can clear all the orders. Alternatively, if he uses only 3 trucks, then all the orders are cleared at the end of the 10th day. What is the minimum number of trucks required so that there will be no pending order at the end of the 5th day?
- (A) 4 (B) 5
(C) 6 (D) 7
64. The variable cost (V) of manufacturing a product varies according to the equation $V = 4q$, where q is the quantity produced. The fixed cost (F) of production of same product reduces with q according to the equation $F = 100/q$. how many units should be produced to minimize the total cost (V+F)?
- (A) 5 (B) 4
(C) 7 (D) 6



Grassroots Academy

